arXiv:math/0701373v1 [math.AP] 13 Jan 2007

A new approach to hyperbolic inverse problems II (Global step)

G.Eskin, Department of Mathematics, UCLA,
Los Angeles, CA 90095-1555, USA. E-mail: eskin@math.ucla.edu

February 2, 2008

Abstract

We study the inverse problem for the second order self-adjoint hyperbolic equation with the boundary data given on a part of the boundary. This paper is the continuation of the author's paper [E].

In [E] we presented the crucial local step of the proof. In this paper we prove the global step. Our method is a modification of the BC-method with some new ideas. In particular, the way of the determination of the metric is new.

1 Introduction.

Let Ω be a bounded domain in $\mathbf{R}^n$, $n \geq 2$, with smooth boundary $\partial\Omega$. Consider the hyperbolic equation of the form:

$$Lu \overset{def}{=} \frac{\partial^2 u}{\partial t^2} + \sum_{j,k=1}^{n} \frac{1}{\sqrt{g(x)}} \left(-i\frac{\partial}{\partial x_j} + A_j(x) \right) \sqrt{g(x)} g^{jk}(x) \left(-i\frac{\partial}{\partial x_k} + A_k(x) \right) u$$

$$+V(x)u = 0 \qquad (1.1)$$

in $\Omega \times (0, T_0)$ with $C^\infty(\overline{\Omega})$ coefficients. Here $\|g^{jk}(x)\|^{-1}$ is the metric tensor, $g(x) = \det \|g^{jk}\|^{-1}$. We assume that

$$u(x,0) = u_t(x,0) = 0 \quad \text{in} \quad \Omega, \qquad (1.2)$$

$$u\,\big|_{\partial\Omega\times(0,T_0)} = f(x,t). \qquad (1.3)$$

Denote by Λ the Dirichlet-to-Neumann (D-to-N) operator, i.e.

$$(1.4)\quad \Lambda f = \sum_{j,k=1}^{n} g^{jk}(x)\left(\frac{\partial u}{\partial x_j} + iA_j(x)u\right)\nu_k \left(\sum_{p,r=1}^{n} g^{pr}(x)\nu_p\nu_r\right)^{-\frac{1}{2}} |_{\partial\Omega\times(0,T_0)},$$

where $\nu = (\nu_1, ..., \nu_n)$ is the unit exterior normal to $\partial\Omega$ with respect to the Euclidean metric. Let Γ_0 be an open subset of $\partial\Omega$. We say that the D-to-N operator is given on $\Gamma_0 \times (0, T_0)$ if $\Lambda f|_{\Gamma_0\times(0,T_0)}$ is known for all smooth $f(x, t)$ with supports in $\Gamma_0 \times (0, T_0]$.

Denote by $G_0(\overline{\Omega})$ the group of all complex-valued functions $c(x)$ such that $c(x) \neq 0$ in $\overline{\Omega}$ and $c(x) = 1$ on $\overline{\Gamma_0}$. We say that potentials $A(x) = (A_1(x), ..., A_n(x))$ and $A'(x) = (A'_1(x), ..., A'_n(x))$ are gauge equivalent if there exists $c(x) \in G_0(\overline{\Omega})$ such that

$$A'_j(x) = A_j(x) - ic^{-1}(x)\frac{\partial c}{\partial x_j}, \qquad 1 \leq j \leq n.$$

Note that if $Lu = 0$ then

$$(1.5)\qquad u' = c^{-1}(x)u$$

satisfies the equation $L'u' = 0$ where L' has the form (1.1) with $A_j(x)$ replaced by $A'_j(x)$, $1 \leq j \leq n$. We shall call (1.5) the gauge transformation.

We shall prove the following theorem:

Theorem 1.1. *Let $L^{(p)}, p = 1, 2$, be two operators of the form (1.1) in domains $\Omega^{(p)}, p = 1, 2$, respectively. Let $\Gamma_0 \subset \partial\Omega^{(1)} \cap \partial\Omega^{(2)}$ and let $\Lambda^{(p)}, p = 1, 2$, be the D-to-N operators corresponding to $L^{(p)}, p = 1, 2$. Assume that $L^{(p)}$ are self-adjoint, i.e. coefficients $A_1^{(1)}(x), ..., A_n^{(1)}(x), V^{(1)}(x)$ and $A_1^{(2)}(x), ..., A_n^{(2)}(x)$, $V^{(2)}(x)$ are real-valued. Suppose $T_0 > 2\max_{x\in\bar{\Omega}^{(1)}} d_1(x, \Gamma_0)$, where $d_1(x, \Gamma_0)$ is the distance in $\overline{\Omega_1}$ with respect to the metric $\|g_1^{jk}(x)\|^{-1}$ from $x \in \overline{\Omega^{(1)}}$ to Γ_0. Suppose that the D-to-N operators $\Lambda^{(1)}$ and $\Lambda^{(2)}$ are equal on $\Gamma_0 \times (0, T_0)$ for all f with supp $f \subset \Gamma_0 \times (0, T_0]$. Then there exists a diffeomorphism φ of $\overline{\Omega_2}$ onto $\overline{\Omega_1}$, $\varphi = I$ on Γ_0, and there exists a gauge transformation $c(x) \in G_0(\overline{\Omega^{(1)}}$ such that $c \circ \varphi \circ L^{(2)} = L^{(1)}$ in $\Omega^{(1)}$.*

An important case of the inverse problems with boundary data on a part of the boundary are the inverse problems in domains with obstacles. In this case $\Omega = \Omega_0 \setminus \cup_{r=1}^{m}\overline{\Omega_r}$, where Ω_0 is diffeomorphic to a ball, $\overline{\Omega_1}, ..., \overline{\Omega_m}$ are smooth nonintersecting domains in Ω_0 called obstacles, $\Gamma_0 = \partial\Omega_0$ and zero Dirichlet boundary conditions hold on $\partial\Omega_r,\ 1 \leq r \leq m$ (c.f. [E1]).

The first result on the inverse problems with the data on a part of the boundary was obtained in [I]. The general self-adjoint case was studied by the BC-method (see [B1], [B2], [K], [KK], [KKL], [KL1]). The present paper is a continuation of the paper [E] (see also [E2]). In [E] the crucial local step was considered, i.e. the unique determination of the coefficients of (1.1) modulo a diffeomorphism and a gauge transformation near Γ_0.

In this paper we complete the proof of Theorem 1.1. In §2 we state the main results proven in [E] and prove the extension lemma. In §3 we refine the results of §2, and in §4 we complete the proof of Theorem 1.1.

2 The summary of the local step and the extension lemma.

Let $L^{(p)}, p = 1, 2$, be two operators of the form (1.1) in $\Omega^{(p)} \times (0, T_0),\ L^{(p)}u_p = 0$ in $\Omega^{(p)} \times (0, T_0),\ u_p(x, 0) = u_{pt}(x, 0) = 0,\ x \in \Omega^{(p)},\ u_p|_{\Gamma_0\times(0,T_0)} = f,\ , p = 1, 2$.

Let Γ be an open connected subset of Γ_0 and let $x = (x', x_n)$ be a system of coordinates in a neighborhood $V \subset \mathbf{R}^n$ of Γ such that $x_n = 0$ is the equation of Γ and $x' = (x_1, ..., x_{n-1})$ are local coordinates on Γ. Introduce semigeodesic coordinates in V corresponding to Γ and to the metric $\|g_p^{jk}\|^{-1}, p = 1, 2$:

$$y = \varphi_p(x). \tag{2.1}$$

Note that $\varphi_p(x) = (\varphi_{p1}(x), ..., \varphi_{pn}(x)), p = 1, 2$, satisfy the following differential equations (see [E], page 817)

$$\sum_{j,k=1}^{n} g_p^{jk}(x)\frac{\partial\varphi_{pn}}{\partial x_j}\frac{\partial\varphi_{pn}}{\partial x_k} = 1,\quad 0 \leq x_n < \delta, \tag{2.2}$$

$$\varphi_{pn}(x', 0) = 0,$$

$$\sum_{j,k=1}^{n} g_p^{jk}(x)\frac{\partial\varphi_{pn}}{\partial x_j}\frac{\partial\varphi_{pr}}{\partial x_k} = 0, \tag{2.3}$$

$$\varphi_{pr}(x', 0) = x_r,\quad 1 \leq r \leq n-1.$$

Denote by $\hat{L}^{(p)})$ the operator $L^{(p)}$ in semigeodesic coordinates:

$$\hat{L}^{(p)}v_p \overset{def}{=} \frac{\partial^2 v_p(y,t)}{\partial t^2} + \frac{1}{\sqrt{\hat{g}_p(y)}}\left(-i\frac{\partial}{\partial y_n} + \hat{A}_{pn}(y)\right)\sqrt{\hat{g}_p(y)}\left(-i\frac{\partial}{\partial y_n} + \hat{A}_{pn}\right)v_p$$
$$+\sum_{j,k=1}^{n-1}\frac{1}{\sqrt{\hat{g}_p(y)}}\left(-i\frac{\partial}{\partial y_j} + \hat{A}_{pj}(y)\right)\sqrt{\hat{g}_p}\hat{g}_p^{jk}(y)\left(-i\frac{\partial}{\partial y_k} + \hat{A}_{pk}(y)\right)v_p(y,t)$$
$$+\hat{V}_p(y)v_p(y,t) = 0,$$

where $v_p(\varphi_p(x),t) = u_p(x,t),\ L^{(p)}u_p = 0,\ p = 1,2$. Denote

$$u_p^{(1)}(y,t) = (\hat{g}_p(y',y_n))^{\frac{1}{4}}e^{-i\psi_p(y',y_n)}u_p(\varphi_p^{-1}(y),t), \quad p = 1,2, \tag{2.4}$$

where the gauge transformation $c_p(y) = e^{-i\psi_p(y)} \in G_0(\overline{U})$ is such that

$$A_{pn}^{(1)}(y) = \hat{A}_{pn}(y) + \frac{\partial\psi_p}{\partial y_n} = 0, \qquad \psi_p(y',0) = 0. \tag{2.5}$$

Then $u_p^{(1)}$ satisfies the equation (c.f. [E], page 819):

$$L_1^{(p)}u_p^{(1)} \overset{def}{=} \frac{\partial^2 u_p^{(1)}}{\partial t^2} - \frac{\partial^2 u_p^{(1)}}{\partial y_n^2} \tag{2.6}$$
$$+\sum_{j,k=1}^{n}\left(-i\frac{\partial}{\partial y_j} + A_{pj}^{(1)}(y)\right)\hat{g}_p^{jk}(y)\left(-i\frac{\partial}{\partial y_k} + A_{pk}^{(1)}(y)\right)u_p^{(1)}$$
$$+V_{p1}u_p^{(1)} = 0, \qquad p = 1,2.$$

Let $T \in (0,T_0)$ be small. As in [E] denote by $\mathcal{D}_p(\Gamma \times (0,T))$ the forward domain of influence of $\overline{\Gamma} \times [0,T]$ for $y_n \geq 0, p = 1,2$, and let $\overline{\mathcal{G}_p} = \{y' : (y',y_n,t) \in \mathcal{D}_p(\Gamma \times (0,T)), y_n = 0, t = T\}$. It follows from the proof of Lemma 2.4 in [E] that $\mathcal{G}_1 = \mathcal{G}_2 \overset{def}{=} \mathcal{G}$ since $\Lambda^{(1)} = \Lambda^{(2)}$ on $\Gamma_0 \times (0,T)$.

Analogously, let $\mathcal{D}_p(\mathcal{G} \times (0,T))$ be the forward domain of influence of $\overline{\mathcal{G}} \times [0,T]$ for $y_n \geq 0, p = 1,2$. Let $Y_{20}^{(p)}$ be the intersection of $\mathcal{D}_p(\mathcal{G} \times (0,T))$ with the plane $T - y_n - t = 0$ and let $X_{20}^{(p)}$ be the part of $\mathcal{D}_p(\mathcal{G} \times (0,T))$ below $Y_{20}^{(p)}, p = 1,2$ (c.f. [E], page 819). Note that $\Lambda^{(1)} = \Lambda^{(2)}$ on $\Gamma_0 \times (0,T_0)$ implies that $\hat{\Lambda}^{(1)} = \hat{\Lambda}^{(2)}$ on $\mathcal{G} \times (0,T)$, where $\hat{\Lambda}^{(p)}f' = \frac{\partial u_p^{(1)}}{\partial y_n}|_{y_n=0}$ is the D-to-N operator corresponding to $L_1^{(p)}$ (see [E], page 819). Here $f' = u_p^{(1)}|_{y_n=0}, u_p^{(1)} = u_{pt}^{(1)} = 0$ for $t = 0, y_n > 0$.

The main result of [E] is the following lemma:

Lemma 2.1. *Let* $T_p, p = 1, 2$, *be such that the semigeodesic coordinates hold in* $X_{20}^{(p)}$, $\overline{\mathcal{G}} \subset \Gamma_0$ *and* $X_{20}^{(p)}$ *does not intersect* $\partial\Omega^{(p)} \times [0, T_p]$ *when* $y_n > 0$. *Suppose* $\hat{\Lambda}^{(1)} = \hat{\Lambda}^{(2)}$ *on* $\mathcal{G} \times (0, T)$, *where* $T = \min(T_1, T_2)$. *Then* $u_1^{(1)} = u_2^{(1)}$ *for* $y' \in \overline{\Gamma}, 0 \leq y_n \leq \frac{T}{2}, y_n \leq t \leq T - y_n$ *and* $L_1^{(1)} = L_1^{(2)}$ *in* $\overline{\Gamma} \times [0, \frac{T}{2}]$.

Remark 2.1 In [E] we assumed that $\mathcal{D}_p(\mathcal{G} \times (0, T_p))$ does not intersect $\partial\Omega^{(p)} \times [0, T_p]$ when $y_n > 0$. It is easy to see that it is enough to assume that $X_{20}^{(p)}$ does not intersect $\partial\Omega^{(p)} \times [0, T_p]$ for $y_n > 0$ since by the domain of dependence argument the solution of $L_1^{(p)} u_p^{(1)} = 0,\ u_p^{(1)} = u_{pt}^{(1)} = 0$ for $t = 0,\ u_p^{(1)} = f'$ for $y_n = 0,\ u_p^{(1)} = 0$ for $(\partial\Omega^{(p)} \times [0, T_p]) \cap \{y_n > 0\}$ restricted to $X_{20}^{(p)}$ does not change whether $(\partial\Omega^{(p)} \times [0, T_p]) \cap \{y_n > 0\}$ intersects $\mathcal{D}(\mathcal{G} \times (0, T_p)) \setminus X_{20}^{(p)}$ or not. $\square$

Denote $D_p = \varphi_p^{-1}(\Gamma \times [0, \frac{T}{2}]) \subset \Omega^{(p)}, p = 1, 2$. Then $\varphi = \varphi_1^{-1} \circ \varphi_2$ is a diffeomorphism of $\overline{D}_2$ onto $\overline{D}_1$. Note that $\hat{g}_1(y', 0) = \hat{g}_2(y', 0)$ (c.f. Remark 2.2 in [E]). It follows from (2.4) that $u_1(\varphi_1^{-1}(y), t) = c_1(y) u_2(\varphi_2^{-1}(y), t)$, where $c_1(y) \in G_0(\overline{\Gamma} \times [0, \frac{T}{2}])$. Therefore

$$c_2 \circ \varphi \circ L^{(2)} = L^{(1)} \text{ in } \overline{D}_1, \tag{2.7}$$

where $c_2(x) = c_1(\varphi_1(x)) \in G_0(\overline{D}_1)$.

Remark 2.2 The subset $\Gamma \subset \Gamma_0$ does not have to be small. There may be no global coordinates $x = (x_1, x_2, ..., x_n)$ near Γ such that $x_n = 0$ is the equation of Γ. In this case we take a finite cover $\{V_j\}, j = 1, ..., N$, of $\overline{\Gamma}$ and apply Lemma 2.1 to each $\Gamma_j = \Gamma \cap V_j,\ j = 1, ..., N$. Let $T_{pj}, \psi_{pj},\ p = 1, 2,\ j = 1, ..., N$, be the same as T_p, φ_p in Lemma 2.1.

Let $T_p = \min_{1 \leq j \leq N} T_{pj},\ T = \min(T_1, T_2)$. Maps $\psi_{pj}, 1 \leq j \leq N$, define a diffeomorphism φ_p of D_p onto the manifold $\overline{\Gamma} \times [0, \frac{T}{2}]$ where $\overline{D}_p \overset{def}{=} \varphi_p^{-1}(\overline{\Gamma} \times [0, \frac{T}{2}]) \subset \overline{\Omega}^{(p)},\ p = 1, 2,\ \varphi_p = I$ on $\overline{\Gamma}$.

Note that sets $\mathcal{G},\ X_{20}^{(p)}$ corresponding to $\Gamma_j,\ 1 \leq j \leq N$ determine sets $\mathcal{G},\ X_{20}^{(p)}$ corresponding to the manifold Γ. Note also that $u_p^{(1)}(y', y_n)$ (see (2.4)) is not a scalar function on $\Gamma \times [0, T]$, but a half-density as $(g_p(y', y_n))^{\frac{1}{4}}$.

Therefore Lemma 2.1 implies:

Lemma 2.2. *Let* Γ *and* T *be such that* $\overline{\mathcal{G}} \subset \Gamma_0$, *the semigeodesic coordinates hold in* $X_{20}^{(p)}, p = 1, 2$, *and* $X_{20}^{(p)} \cap (\partial\Omega^{(p)} \setminus (\Gamma_0 \cap \cup_{j=1}^N V_j)) = \emptyset$. *Suppose* $\hat{\Lambda}^{(1)} = \hat{\Lambda}^{(2)}$ *on* $\mathcal{G} \times (0, T)$. *Denote* $\varphi = \varphi_1^{-1} \circ \varphi_2$. *There exists a gauge* *transformation* $c_2 \in G_0(\overline{D}_1)$ *such that (2.7) holds. or equivalently*

$$L_{1j}^{(1)} = L_{1j}^{(2)} \quad in \quad \overline{\Gamma}_j \times [0, \frac{T}{2}], \quad 1 \le j \le N, \tag{2.8}$$

where $L_{1j}^{(p)}, p = 1, 2,$ *have the form (2.6) in local semigeodesic coordinates in* $\overline{\Gamma}_j \times [0, \frac{T}{2}]$. *For the brevity of notations we shall write* $L_1^{(1)} = L_1^{(2)}$ *in* $\overline{\Gamma} \times [0, \frac{T}{2}]$ *instead of (2.8).*

Note that $\overline{D}_p$ is the union of all geodesics in $\overline{\Omega}^{(p)}, p = 1, 2,$ starting at $\overline{\Gamma}$, orthogonal to $\overline{\Gamma}$ and having the length $\frac{T}{2}$.

Remark 2.3 In §4 we shall deal often with the following situation: Let V be the same neighborhood as in the beginning of this section. Suppose $L^{(p)}$ are defined also in $\overline{V}$, $p = 1, 2$, and $L^{(1)} = L^{(2)}$ when $x_n < 0$, i.e. in $V_- \overset{def}{=} V \setminus \overline{\Omega}^{(p)}$. Let $y = \varphi_p(x)$ be the same as in (2.2), (2.3). Therefore $\varphi = \varphi_1^{-1} \circ \varphi_2$ is the diffeomorphism of $\overline{D}_2$ onto $\overline{D}_1$. We shall show that defining $\varphi = I$ on V_- we get a diffeomorphism of $\overline{D}_2 \cup \overline{V}_-$ onto $\overline{D}_1 \cup \overline{V}_-$.

It follows from (2.2), (2.3) and (2.7) that $\varphi(x)$ satisfies the equations:

$$\|g_1^{jk}\| = \frac{D\varphi}{Dx} \, \|g_2^{jk}\| \left(\frac{D\varphi}{Dx}\right)^T \quad \text{in} \quad V,$$

where $\frac{D\varphi}{Dx}$ is the Jacobi matrix of φ. Since $\varphi = I$ when $x_n = 0$ and since $\|g_1^{jk}\| = \|g_2^{jk}\|$ when $x_n \le 0$ we have that $\varphi = I$ for $x_n \le 0$. □

Let $B \subset \Gamma \times [0, \frac{T}{2}]$ be a domain such that $\partial B \cap \Gamma$ is open and connected. Denote $B_p = \varphi_p^{-1}(B)$. We assume that $\Omega^{(p)} \setminus \overline{B}_p$ is smooth. Note that the restriction of $\varphi = \varphi_1^{-1} \circ \varphi_2$ to $\overline{B}_2$ maps $\overline{B}_2$ onto $\overline{B}_1$. The following extension lemma holds (see [Hi], Chapter 8):

Lemma 2.3. *There exists a diffeomorphism* φ_3 *of* $\overline{\Omega}^{(2)}$ *onto* $\overline{\Omega}^{(3)} \overset{def}{=} \varphi_3(\overline{\Omega}^{(2)})$ *such that* $\varphi_3|_{B_2} = \varphi$ *and* $\varphi_3|_{\Gamma_0} = I$.

Proof: Since $\overline{D}_2 = \varphi_2^{-1}(\overline{\Gamma} \times [0, \frac{T}{2}])$ and $\varphi = \varphi_1 = \varphi_2 = I$ on $\overline{\Gamma}$ there exists a smooth family $\psi_t, 0 \le t \le 1$, of embedding of $\overline{D}_2$ in $\mathbf{R}^n$ such that $\psi_0 = \varphi$ and $\psi_1 = I$ on $\overline{D}_2$. The proof of this fact is similar to the proof of Theorem 8.3.1. in [Hi]. Denote $A = \overline{B}_2 \cup \overline{\Gamma}_0$. Define $\varphi = I$ on $\overline{\Gamma}_0$. Denote by B_2' the union of a small neighborhood of $\overline{D}_2$ and a small neighborhood of $\overline{\Gamma}_0$. We can extend $\psi_t, 0 \le t \le 1$ from $\overline{D}_2$ to B_2' preserving the properties that $\psi_0 = \varphi$ on A, $\psi_t = I$ on $\overline{\Gamma}_0$, $\psi_1 = I$ on B_2', ψ_t is an embedding of B_2' in $\mathbf{R}^n$. Now applying Theorem 8.1.4 in [Hi] we get that there exists a diffeomorphism φ_3 of $\mathbf{R}^n$ onto $\mathbf{R}^n$, $\varphi_3 = I$ for $|x| > N$, N is large, such that $\varphi_3|_A = \varphi$. Taking the restriction of φ_3 to $\overline{\Omega}^{(2)}$ we prove Lemma 2.3. $\square$

Denote by $c_3 \in G_0(\overline{\Omega}^{(3)})$ the extension of c_2 from $\overline{B}_1 \cup \overline{\Gamma}_0$ to $\overline{\Omega}^{(3)}$, where c_2 is the same as in (2.7), $c_2 = 1$ on Γ_0 , $\Omega^{(3)} \overset{def}{=} \varphi_3(\Omega^{(2)})$. Let $L^{(3)} = c_3 \circ \varphi_3 \circ L^{(2)}$ be the differential operator in $\Omega^{(3)}$. We have that $L^{(3)} = L^{(1)}$ in $\overline{B}_1$. Note that $B_1 \subset \Omega^{(1)} \cap \Omega^{(3)}$ and $\Gamma_0 \subset \partial\Omega^{(1)} \cap \partial\Omega^{(3)}$.

Let $\Gamma_1 = (\Gamma_0 \setminus (\partial B_1 \cap \Gamma_0)) \cup (\partial B_1 \setminus \overline{\Gamma}_0)$ and let $\delta = \max_{x\in\overline{B}_1} d_1(x, \Gamma_0)$, where $d_1(x, \Gamma_0)$ is the distance in $\overline{B}_1$ from $x \in \overline{B}_1$ to Γ_0. Note that $\Gamma_1 \subset \partial(\Omega^{(1)} \setminus \overline{B}_1) \cap \partial(\Omega^{(3)} \setminus \overline{B}_1)$.

Let $\Lambda^{(1)}$ and $\Lambda^{(3)}$ be the D-to-N operators corresponding to $L^{(1)}$ and $L^{(3)}$ in domains $\Omega^{(1)} \times (0, T_0)$ and $\Omega^{(3)} \times (0, T_0)$, respectively. For the simplicity of notations we continue to denote by $\Lambda^{(1)}$, $\Lambda^{(3)}$ the D-to-N operators corresponding to $L^{(1)}$ and Ł$^{(3)}$ in smaller domains $(\Omega^{(1)} \setminus \overline{B}_1) \times (\delta, T_0 - \delta)$, $(\Omega^{(3)} \setminus \overline{B}_1) \times (\delta, T_0 - \delta)$, respectively.

The following lemma was proven in [E], Lemma 3.3.

Lemma 2.4. *If* $\Lambda^{(1)} = \Lambda^{(3)}$ *on* $\Gamma_0 \times (0, T_0)$ *then the D-to-N operators corresponding to* $L^{(1)}$, $L^{(3)}$ *on domains* $(\Omega^{(1)} \setminus \overline{B}_1) \times (\delta, T_0 - \delta)$, $(\Omega^{(3)} \setminus \overline{B}_1) \times (\delta, T_0 - \delta)$ *respectively are equal on* $\Gamma_1 \times (\delta, T_0 - \delta)$.

Note that the inverse of Lemma 2.4 is also true:

Lemma 2.5. *Suppose the D-to-N operators corresponding to* $L^{(1)}$, $L^{(3)}$ *on domains* $(\Omega^{(1)} \setminus \overline{B}_1) \times (0, T_0)$, $(\Omega^{(1)} \setminus \overline{B}_1 / \times (0, T_0)$, *respectively, are equal on* $\Gamma_1 \times (0, T_0)$. *Then* $\Lambda^{(1)} = \Lambda^{(3)}$ *on* $\Gamma_0 \times (0, T_0)$.

Proof: Let f be a smooth function on $\partial\Omega^{(1)} \times (0, T_0)$, $\operatorname{supp} f \subset \Gamma_0 \times (0, T_0]$. Let u_1 be the solution of the initial-boundary value problem

$$\begin{gathered} L^{(1)} u_1 = 0 \quad \text{in} \quad \Omega^{(1)} \times (0, T_0), \\ u_1(x, 0) = u_{1t}(x, 0) = 0, \quad x \in \Omega^{(1)}, \qquad u_1|_{\partial\Omega^{(1)} \times (0, T_0)} = f. \end{gathered}$$

Denote by f_1 the restriction of u_1 to $\Gamma_1 \times (0, T_0)$. Let u_3 be the solution of

$$\begin{aligned} L^{(3)} u_3 = 0 \quad \text{in} \quad (\Omega^{(3)} \setminus \overline{B}_1) \times (0, T_0), \\ u_3(x, 0) = u_{3t}(x, 0) = 0, \quad x \in \Omega^{(3)} \setminus \overline{B}_1, \\ u_3|_{\Gamma_1 \times (0, T_0)} = f_1, \quad u_3 = 0 \ \text{ on } \quad ((\partial(\Omega^{(3)} \setminus \overline{B}_1)) \setminus \overline{\Gamma}_1) \times (0, T_0). \end{aligned}$$

Note that u_1 is also zero on $((\partial(\Omega^{(1)} \setminus \overline{B_1})) \setminus \overline{\Gamma_1}) \times (0, T_0)$. We have that $\Lambda^{(1)} f_1 = \Lambda^{(3)} f_1$ on $\Gamma_1 \times (0, T_0)$ by the assumption.

Let $\tilde{u}_3 = u_3$ in $(\Omega^{(3)} \setminus \overline{B}_1) \times (0, T_0)$ and $\tilde{u}_3 = u_1$ in $\overline{B}_1 \times (0, T_0)$. Since $\Lambda^{(1)} f_1 = \Lambda^{(3)} f_1$ and $L^{(3)} = L^{(1)}$ in $\overline{B_1}$ we get that $\tilde{u}_3$ is the solution of $L^{(3)} \tilde{u}_3 = 0$ in $\Omega^{(3)} \times (0, T_0)$. Note that $\tilde{u}_3(x, 0) = \tilde{u}_{3t}(x, 0) = 0$ for $x \in \Omega^{(3)}$. Also $\tilde{u}_3|_{\Gamma_0 \times (0,T_0)} = u_1|_{\Gamma_0 \times (0,T_0)} = f$, $\tilde{u}_3 = 0$ on $(\partial\Omega^{(3)} \setminus \Gamma_0) \times (0, T_0)$. Moreover, $\Lambda^{(3)} f = \Lambda^{(1)} f$ on $\Gamma_0 \times (0, T_0)$ since $u_1 = u_3$ in $\overline{B_1} \times (0, T_0)$ and $\Lambda^{(1)} = \Lambda^{(3)}$ on $\Gamma_1 \times (0, T_0)$. $\square$

3 Refinements of Lemma 2.2.

In this and the next sections we shall show how to use repeatedly Lemmas 2.1 - 2.5 to prove Theorem 1.1. We shall prove the following lemma considering, for the simplicity, first the case when $\Gamma = \Gamma_0$ and $\partial\Gamma_0 = \emptyset$.

Lemma 3.1. *Let $\partial\Gamma_0 = \emptyset$ and let T_1 be such that the semigeodesic coordinates for $L^{(1)}$ hold in $D_1 = \varphi_1^{-1}(\Gamma_0 \times [0, \frac{T_1}{2}])$ and $D_1 \cap (\partial\Omega^{(1)} \setminus \Gamma_0) = \emptyset$. Let T_2^* be such that the semigeodesic coordinates for $L^{(2)}$ hold in $\overline{D}_2 = \varphi_2^{-1}(\Gamma_0 \times [0, \frac{T_2}{2}])$ for any $T_2 < T_2^*$ and $D_2 \cap (\partial\Omega^{(2)} \setminus \Gamma_0) = \emptyset$ for any $T_2 < T_2^*$. Suppose that there exists a focal point of a geodesics γ_0 in $\overline{\Omega}^{(2)}$, starting at Γ_0, orthogonal to Γ_0 and such that $y_{n0} = \frac{T_2^*}{2}$, where (y_0', y_{n0}) are semigeodesic coordinates of the focal point. Then, assuming that $\Lambda^{(1)} = \Lambda^{(2)}$ on $\Gamma_0 \times (0, T_0)$ and $T_0 > T_1$, we have that $T_2^* > T_1$ and $L_1^{(1)} = L_1^{(2)}$ in $\Gamma_0 \times [0, \frac{T_1}{2}]$.*

Proof: Consider the bicharacteristics system:

$$
\begin{aligned}
&\frac{dx_{rj}}{dt} = \frac{\partial H_r(x_r, p_r)}{\partial p_j}, \quad x_{rj}(0) = y_j, \ 1 \le j \le n-1, \ r = 1, 2, \\
&\frac{dx_{rn}}{dt} = \frac{\partial H_r(x_r, p_r)}{\partial p_n}, \quad x_{rn}(0) = 0, \quad r = 1, 2, \\
&\frac{dp_{rj}}{dt} = -\frac{\partial H_r(x_r, p_r)}{\partial x_j}, \quad p_{rj}(0) = 0, \ 1 \le j \le n-1, \ r = 1, 2, \\
&\frac{dp_{rn}}{dt} = -\frac{\partial H_r(x_r, p_r)}{\partial x_n}, \quad p_n(0) = \frac{1}{\sqrt{g_r^{nn}(y', 0)}}, \quad r = 1, 2.
\end{aligned}
\tag{3.1}
$$

Here $H_r(x, p) = \sqrt{\sum_{j,k=1}^n g_r^{jk}(x) p_j p_k}$. Let $\varphi_r(x) = (\varphi_{r1}(x), ..., \varphi_{rn}(x))$ be the same as in (2.2), (2.3). Then $p_{rj}(t) = \frac{\partial\varphi_{rn}(x_r(t))}{\partial x_j}$, $1 \le j \le n$, $r = 1, 2$, and

(2.2) implies that $\frac{d\varphi_{rn}(x(t))}{dt} = \sum_{j=1}^n p_{rj}(t)\frac{\partial H_r(x_r,p_r)}{\partial p_j} = H(x_r(t), p_r(t)) = 1$.

Since $\varphi_{rn}(x(0)) = 0$ we get that

$$\varphi_{rn}(x_r(t)) = t. \tag{3.2}$$

Note that (2.3) implies:

$$\frac{d\varphi_{rj}(x_r(t))}{dt} = \sum_{k=1}^n \frac{\partial\varphi_{rj}(x_r(t))}{\partial x_k}\frac{dx_{rk}}{dt} = \sum_{k=1}^n \frac{\partial\varphi_{rj}(x_r(t))}{\partial x_k}\frac{\partial H_r(x_r, \frac{\partial\varphi_{rn}}{\partial x})}{\partial p_k} = 0,$$
$$1 \le j \le n-1.$$

Therefore

$$\varphi_{rj}(x_r(t)) = \varphi_{rj}(x_r(0)) = \varphi_{rj}(y', 0) = y_j,\ 1 \le j \le n-1,\ r = 1, 2. \tag{3.3}$$

Denote $y_n = t$. The coordinates $(y_1, ..., y_{n-1}, y_n)$ are the semigeodesic coordinates. Consider the change of variables

$$x_j = x_{rj}(y_1, ..., y_n), \quad 1 \le j \le n, \quad r = 1, 2. \tag{3.4}$$

Assume that the Jacobian

$$\det\frac{D(x_{r1}, ..., x_{rn})}{D(y_1, ..., y_n)} \ne 0.$$

It follows from (3.2), (3.3) that (3.4) is the inverse to the change of variables (2.1), since we have :

$$y_j = \varphi_{rj}(x_r(y_1, ..., y_{n-1}, t)),$$
$$y_n = \varphi_{rn}(x_r(y_1, ..., y_{n-1}, t)) = t.$$

Therefore $\frac{D\varphi_r(x)}{Dx} = (\frac{Dx_r}{Dy})^{-1}$. Note that the solution of the system (3.1) exists for all $t \in \mathbf{R}$. Therefore $\frac{Dx_r}{Dy}$ is smooth for all y. However the Jacobian $\det\frac{Dx_r}{Dy}$ may vanish for some y. Such y are called the focal points (caustics) for the Hamiltonian system (3.1). Then $\det\frac{D\varphi_r}{Dx} \to \infty$ when $y = \varphi_r(x)$ approaches a focal point. Suppose $T_2^* \le T_1$. Then there exists a focal point $y^{(0)} = (y_0', y_{n0}) \in \Gamma_0 \times [0, \frac{T_1}{2}]$ for the system (3.1) when $r = 2$ and $y_{n0} \le \frac{T_1}{2}$. We have near $y^{(0)}$(see (1.8) in [E]):

$$\det\|\hat{g}_r^{jk}(\varphi_r(x))\| = \det\|g_r^{jk}(x)\|\left(\det\frac{D\varphi_r}{Dx}\right)^2, \tag{3.5}$$

where $\|\hat{g}_r^{jk}(y)\|^{-1}$ is the metric tensor in semigeodesic coordinates. If $y \in \Gamma_0 \times [0, \frac{T_2}{2}]$ and $\frac{T_2}{2}$ is close to y_{n0}, $\frac{T_2}{2} < y_{n0} \leq \frac{T_1}{2}$, we have, by Lemma 2.2 that $\hat{g}_1^{jk} = \hat{g}_2^{jk}$. Consider (3.5) when $r = 1$. Since $y^{(0)}$ is not a focal point for $r = 1$ we have that $\det \frac{D\varphi_1}{Dx}$ is bounded. Therefore $\det \|\hat{g}_1^{jk}\|$ is bounded near $y^{(0)}$. Now consider (3.5) for $r = 2$. Then $\left(\det \frac{D\varphi_2}{Dx}\right)^2 = \frac{\det \|\hat{g}_1^{jk}\|}{\det \|g_2^{jk}\|}$ is also bounded when $y \to y^{(0)}$. Therefore $y^{(0)}$ is not a focal point for $r = 2$ and this is a contradiction. Therefore we can take $T_2 = T_1$ and we have $L_1^{(1)} = L_1^{(2)}$ in $\Gamma_0 \times [0, \frac{T_1}{2}]$ (c.f. (2.8)). □

Lemma 3.2. *Let $\partial\Gamma_0 = \emptyset$ and let T_1 be such that the semigeodesic coordinates for $L^{(1)}$ hold in $D_1 = \varphi_1^{-1}(\Gamma_0 \times [0, \frac{T_1}{2}])$ and $D_1 \cap (\partial\Omega^{(1)} \setminus \Gamma_0) = \emptyset$. Then semigeodesic coordinates for $L^{(2)}$ hold also in $D_2 = \varphi_2^{-1}(\Gamma_0 \times [0, \frac{T_1}{2}])$, $D_2 \cap (\partial\Omega^{(2)} \setminus \Gamma_0) = \emptyset$ and $L_1^{(1)} = L_1^{(2)}$ in $\Gamma_0 \times [0, \frac{T_1}{2}]$.*

Proof: Suppose D_2 intersects $\partial\Omega^{(2)} \setminus \Gamma_0$. Let $x^{(0)}$ be the point in $D_2 \cap (\partial\Omega^{(2)} \setminus \Gamma_0)$ closest to Γ_0 and let $y^{(0)} = (y_0', y_{n0})$ be the semigeodesic coordinates of $x^{(0)}$. We have by assumption that $y_{n0} \leq \frac{T_1}{2}$. For any $\frac{T_2}{2} < y_{n0}$ we have by Lemma 3.1 that $L_1^{(1)} = L_1^{(2)}$ in $\Gamma_0 \times [0, \frac{T_2}{2}]$. Therefore by the continuity $L_1^{(1)} = L_1^{(2)}$ in $\Gamma_0 \times [0, y_{n0}]$.

Denote by γ_0 the geodesics in $\overline{\Omega}^{(2)}$ starting at $x^{(1)} \in \Gamma_0$, orthogonal to Γ_0 and reflecting at $\partial\Omega^{(2)} \setminus \Gamma_0$ at point $x^{(0)}$. Since $x^{(0)}$ is the closest point to Γ_0 in D_1, the angle of reflection at $x^{(0)}$ is $\frac{\pi}{2}$. Denote by u_2 the geometric optics solution depending on large parameter associated with γ_0 and its successive reflections (c.f. [E], page 824, and [E2], pages 28-29). We have $L^{(2)}u_2 = 0$ in $\Omega^{(2)} \times (0, T_0)$, $u_2 = u_{2t} = 0$ for $t = 0$, and $u_2|_{\partial\Omega^{(2)} \times (0,T_0)} = f$ where supp f is contained in a small neighborhood of $(x^{(1)}, t_1) \in \Gamma_0 \times (0, T_0), t_1 > 0$, where $(x^{(1)}, t_1)$ is the starting point of the broken ray γ_0. In semigeodesic coordinates u_2 is concentrated (modulo lower order terms) in a small neighborhood of a broken ray $y' = y_0'$, $y_n - t = -t_1$ for $t_1 < t < y_{n0} + t_1$, $y' = y_0'$, $y_n + t = 2y_{n0} + t_1$ for $y_{n0} + t_1 < t < t_1 + 2y_{n0}$, $y' = y_0'$, $y_n - t = -t_1 - 2y_{n0}$ for $t > t_1 + 2y_{n0}$, etc.

Denote by u_1 the solution of $L^{(1)}u_1 = 0$ in $\Omega^{(1)} \times (0, T_0)$, $u_1 = u_{1t} = 0$ for $t = 0$, $u_1|_{\partial\Omega^{(1)} \times (0,T_0)} = f$. Let $u_p^{(1)}$ be related to $u_p, p = 1, 2$, by formulas (2.4). Since $\Lambda^{(1)} = \Lambda^{(2)}$ on $\Gamma_0 \times (0, T_0)$ we have that $u_1^{(1)}$ and $u_2^{(1)}$ have tha same Cauchy data when $y_n = 0$. Since $L_1^{(1)} = L_1^{(2)}$ in $\Gamma_0 \times [0, y_{n0}]$ we get, by the unique continuation theorem (see [T]), that $u_1^{(1)} = u_2^{(1)}$ for

$y \in \Gamma_0 \times [0, y_{n0}],\ 0 < t < T_0 - y_{n0}$. However, $y^{(0)}$ is not a point of reflection for $u_1^{(1)}$. Therefore $u_1^{(1)} \neq u_2^{(1)}$ for $y_{n0} + t_1 < t < y_{n0} + t_1 + \varepsilon,\ y_n < y_{n0}$, where $\varepsilon > 0$ is small. This contradiction proves that $(\partial\Omega^{(2)} \setminus \Gamma_0) \cap \bar{D}_2 = \emptyset$. $\square$

Now we shall consider the case when $\Gamma \subset \Gamma_0$ may have a boundary, i.e. when $\partial\Gamma \neq \emptyset$. Let T_1 be such that all conditions of Lemma 2.2 are satisfied. Let $X_{j0}^{(p)}, j = 1, 2$, be the same as in Lemma 2.2. Denote by $\Delta_j^{(p)}$ the projection of $X_{j0}^{(p)}$ on the plane $t = 0, j = 1, 2, p = 1, 2$. Note that $\Delta_j^{(p)}$ are contained in the strip $0 \leq y_n \leq \frac{T_1}{2}$. Let $\tilde{\Gamma} \subset \Gamma$ be such that $\overline{\tilde{\Delta}}_2^{(1)} \subset \Gamma \times [0, \frac{T_1}{2}]$ where $\tilde{\Delta}_2^{(p)}$ is the same as $\Delta_2^{(p)}$ when Γ is replaced by $\tilde{\Gamma}$.

We will need the following proposition:

Lemma 3.3. *Let $L^{(p)}, p = 1, 2$, be two operators such that $L_1^{(1)} = L_1^{(2)}$ in $(\overline{\Gamma} \setminus \tilde{\Gamma}) \times [0, \frac{T_1}{2}]$. Suppose $\overline{\tilde{\Delta}}_2^{(1)} \subset \Gamma \times [0, \frac{T_1}{2}]$. Then $\overline{\tilde{\Delta}}_2^{(2)} \subset \overline{\tilde{\Delta}}_2^{(1)}$.*

Proof: Suppose there exists $(y_0', y_{n0}) \in \overline{\tilde{\Delta}}_2^{(2)}$, where $y_{n0} \leq \frac{T_1}{2},\ (y_0', y_{n0}) \notin \overline{\tilde{\Delta}}_2^{(1)}$. Note that $(y_0', y_{n0}) \in \overline{\tilde{\Delta}}_2^{(2)}$ means that there exists $y_1' \in \overline{\tilde{\mathcal{G}}}$ such that the shortest path γ connecting $(y_1', 0)$ and (y_0', y_{n0}) has the length $|\gamma| \leq T_1 - y_{n0}$. Here $\tilde{\mathcal{G}}$ is the same for $\tilde{\Gamma}$ as $\mathcal{G}$ for Γ (see §2). If γ is contained completely in $(\overline{\Gamma} \setminus \tilde{\mathcal{G}}) \times [0, \frac{T_1}{2}]$ then we must have $(y_0', y_{n0}) \in \overline{\tilde{\Delta}}_2^{(1)}$ since $L_1^{(1)} = L_1^{(2)}$ in $(\overline{\Gamma} \setminus \tilde{\Gamma}) \times [0, \frac{T_1}{2}]$. Suppose there is a part of γ that does not belong to $(\overline{\Gamma} \setminus \tilde{\mathcal{G}}) \times [0, \frac{T_1}{2}]$. Let $(y_2', y_{n2}) \in \partial\tilde{\mathcal{G}} \times [0, \frac{T_1}{2}]$ be such that the part γ_1 of γ connecting (y_2', y_{n2}) and (y_0', y_{n0}) is in $(\overline{\Gamma} \setminus \tilde{\mathcal{G}}) \times [0, \frac{T_1}{2}]$. Denote by γ_2 the remaining part of γ. Let γ_2' be the straight line connecting (y_2', y_{n2}) and $(y_2', 0)$. Since the metric for $L_1^{(2)}$ has the form $(dy_n)^2 + \sum_{j,k=1}^{n-1} \hat{g}_{2jk} dy_j dy_k$ we have that the length $|\gamma_2'|$ of γ_2' is less or equal than $|\gamma_2|$. Therefore $T - y_{n0} \geq |\gamma| = |\gamma_1| + |\gamma_2| \geq |\gamma_1| + |\gamma_2'|$. The path $\gamma_2' \cup \gamma_1$ is contained in $(\bar{\Gamma} \setminus \tilde{\mathcal{G}}) \times [0, \frac{T_1}{2}]$. Since $L_1^{(1)} = L_1^{(2)}$ in $(\bar{\Gamma} \setminus \tilde{\Gamma}) \times [0, \frac{T_1}{2}]$ we have that $\gamma_2' \cup \gamma_1$ is contained in $\overline{\tilde{\Delta}}_2^{(1)}$. Therefore $(y_0', y_{n0}) \in \overline{\tilde{\Delta}}_2^{(1)}$. This contradiction proves Lemma 3.3. $\square$

The following lemma generalizes Lemmas 3.1 and 3.2 to the case when $\partial\Gamma \neq \emptyset$.

Lemma 3.4. *Consider $\Gamma \subset \Gamma_0$ such that $\partial\Gamma \neq \emptyset$. Let $0 < T_1 < T_0$ be such that the semigeodesic coordinates for $L^{(1)}$ hold in $\varphi_1^{-1}(\bar{\Gamma} \times [0, \frac{T_1}{2}])$ and $(\partial\Omega^{(1)} \setminus \Gamma) \cap \varphi_1^{-1}(\bar{\Gamma} \times [0, \frac{T_1}{2}]) = \emptyset$. Let $\tilde{\Gamma} \subset \Gamma$ be such that $\overline{\tilde{\Delta}_2^{(1)}} \subset \Gamma \times [0, \frac{T_1}{2}]$.*

Suppose that semigeodesic coordinates for $L^{(2)}$ *hold in* $(\bar{\Gamma}\setminus\tilde{\Gamma})\times[0,\frac{T_1}{2}]$, $(\partial\Omega^{(2)}\setminus\Gamma)\cap\varphi_2^{-1}((\bar{\Gamma}\setminus\tilde{\Gamma})\times[0,\frac{T_1}{2}])=\emptyset$ *and* $L_1^{(1)}=L_1^{(2)}$ *in* $(\bar{\Gamma}\setminus\tilde{\Gamma})\times[0,\frac{T_1}{2}]$. *Then the semigeodesic coordinates for* $L^{(2)}$ *hold in* $\bar{\Gamma}\times[0,\frac{T_1}{2}]$, $(\partial\Omega^{(2)}\setminus\Gamma)\cap\varphi_2^{-1}(\bar{\Gamma}\times[0,\frac{T_1}{2}])=\emptyset$ *and* $L_1^{(1)}=L_1^{(2)}$ *in* $\bar{\Gamma}\times[0,\frac{T_1}{2}]$.

Proof: Since $\overline{\tilde{\Delta}}_2^{(2)}\subset\overline{\tilde{\Delta}}_2^{(1)}\subset\Gamma\times[0,\frac{T_1}{2}]$ (see Lemma 3.3) we have that there is no focal points for $L_1^{(2)}$ in $\tilde{\Delta}_2^{(2)}\setminus(\tilde{\Gamma}\times[0,\frac{T_1}{2}])$. Repeating the proof of Lemma 3.1 with Γ_0 replaced by $\tilde{\Gamma}$ we get that there is no focal points for $L_1^{(2)}$ in $\overline{\tilde{\Gamma}}\times[0,\frac{T_1}{2}]$ and $L_1^{(1)}=L_1^{(2)}$ in $\overline{\tilde{\Gamma}}\times[0,\frac{T_1}{2}]$ assuming that $Z\overset{def}{=}(\partial\Omega^{(2)}\setminus\Gamma)\cap\varphi_2^{-1}(\overline{\tilde{\Gamma}}\times[0,\frac{T_1}{2}])=\emptyset$. Now we shall show that the set Z is empty. Suppose $Z\neq\emptyset$. Since $(\partial\Omega^{(2)}\setminus\Gamma)\cap\varphi_2^{-1}(\partial\tilde{\Gamma}\times[0,\frac{T_1}{2}])=\emptyset$ the point $(y'_{n0},y_{n0})\in Z$ closest to Γ belongs to $\tilde{\Gamma}\times[0,\frac{T_1}{2}]$, i.e. y'_0 is an interior point of $\tilde{\Gamma}$.

From this point we can repeat the proof of Lemma 3.2 to get a contradiction. □

4 The global step.

We start this section with a lemma (Lemma 4.1) that will play a key role in the global step of the proof of Theorem 1.1

Let $O(\delta_1)$ be a ball in Γ_0 : $O(\delta_1)=\{x'\in\Gamma_0,\ d_1(x',x'_0)<\delta_1\}$, where $x'_0\in\Gamma_0,\ d_1(x',x'_0)$ is the distance on Γ_0 induced by the metric $\|g_1^{jk}\|^{-1}$. Let R be the union of geodesics in $\Omega^{(1)}$ with respect to the metric $\|g_1^{jk}\|^{-1}$, starting on $\overline{O}(\delta_1)$, orthogonal to Γ_0 and having the lengths $T,\ 2T<T_0$. We assume that these geodesics have no focal points in $\overline{R}$ and do not intersect $\partial\Omega^{(1)}\setminus\overline{O}(\delta_1)$. Therefore we can introduce semigeodesic coordinates $y=\psi_1(x)$ in $\overline{R}$ using these geodesics. By the continuity the semigeodesic coordinates $y=\psi_1(x)$ hold in a larger domain $R_0\supset R$, i.e. there exists a small $\delta_2>0$ such that $\overline{O}(\delta_1+\delta_2)\subset\Gamma_0$ and $y=\psi_1(x)$ is a diffeomorphism of $\overline{R}_0$ onto $\overline{O}(\delta_1+\delta_2)\times[0,T]$.

Denote by R_1 the union of all geodesics in R_0 with the lengths δ_3. We will choose $\delta_3<\delta_2$. Let R'_1 be the union of geodesics in $\Omega^{(2)}$ with respect to the metric $\|g_2^{jk}\|^{-1}$ starting on $\overline{O}(\delta_1+\delta_2)$ orthogonal to Γ_0 and having the lenghts δ_3. Let $y=\psi_2(x)$ be the semigeodesic coordinates on R'_1, i.e. ψ_2 is a diffeomorphism of $\overline{R'_1}$ onto $\overline{O}(\delta_1+\delta_2)\times[0,\delta_3]$.



Figure 1. Domains R_k and boundaries $\hat{\Sigma}_{k-1}$, $k \geq 2$. $\hat{\Sigma}_{k-1}$ is drawn in bold.

Note that the metrics on Γ_0 induced by $\|g_1^{jk}\|^{-1}$ and $\|g_2^{jk}\|^{-1}$ are the same since $\Lambda^{(1)} = \Lambda^{(2)}$ on $\Gamma_0 \times (0, T_0)$ (c.f., for example, Remark 2.2 in [E]). Note that $\psi_3 = \psi_1^{-1} \circ \psi_2$ is a diffeomorphism of $\overline{R}_1'$ onto $\overline{R}_1$. Since $\delta_3 > 0$ is small we can apply Lemma 2.1 or Lemma 2.2 to get $L^{(1)} = L^{(3)}$ in $\overline{R}_1$ where $L^{(3)} = c_3 \circ \psi_3 \circ L^{(2)},\ c_3 \in G_0(\overline{R}_1),\ \psi_3 = I$ and $c_3 = 1$ on $O(\delta_1 + \delta_2),\ L^{(3)}$ is an operator on $\overline{R}_1$.

Let $\hat{X}_{20}^{(p)}$ be the same as in §2 when Γ is replaced by $O(\delta_1 + \frac{\delta_2}{2})$ and let $\hat{\Delta}_{20}^{(p)}$ be the projection of $\hat{X}_{20}^{(p)}$ on the plane $t = 0, p = 1, 3$. If δ_3 is much smaller than δ_2, we have that $\overline{\hat{\Delta}}_{20}^{(1)} \subset O(\delta_1 + \delta_2) \times [0, \delta_3]$ and therefore $\overline{\tilde{\Delta}}_{20}^{(3)} \subset \overline{\tilde{\Delta}}_{20}^{(1)}$ (c.f. Lemma 3.3).

Let $\Sigma_1 = \psi_1^{-1}(\sigma_1)$ where $\sigma_1 = \overline{O}(\delta_1 + \delta_2) \setminus O(\delta_1)$ when $y_n = 0,\ \sigma_1 = \partial O(\delta_1)$ when $0 \leq y_n \leq \delta_3,\ \sigma_1 = O(\delta_1)$ when $y_n = \delta_3$ (see Fig. 1). Here $y = \psi_1(x)$ are the semigeodesic coordinates in $\overline{R}_0$. Note that σ_1 has edges when $y_n = 0,\ y' \in \partial O(\delta_1)$ and when $y_n = \delta_3,\ y' \in \partial O(\delta_1)$. We will smooth $\Sigma_1 = \psi_1^{-1}(\sigma_1)$ near these edges to obtain a smooth surface Σ_1. We can arrange the smoothing in such a way that Σ_1 and $\hat{\Sigma}_1$ differ only in a small neighborhood of edges of the size $O(\varepsilon)$ where $0 < \varepsilon \ll \delta_3$. Denote by R_2 the domain bounded by $\hat{\Sigma}_1$ and $O(\delta_1 + \delta_2)$. Note that $R_2 \subset R_1$. Using Lemma 2.3 we can extend ψ_3 from $\overline{R}_2' \overset{def}{=} \psi_3^{-1}(\overline{R}_2) \subset \overline{\Omega}^{(2)}$ to $\overline{\Omega}^{(2)}$ as a diffeomorphism of $\overline{\Omega}^{(2)}$ onto $\overline{\Omega}^{(3)} \overset{def}{=} \psi_3(\Omega^{(2)})$, $\psi_3 = I$ on Γ_0 and extend c_3 from $\overline{R}_2$ to $\Omega^{(3)}$ as an element of $G_0(\overline{\Omega}^{(3)})$. It follows from Lemma 2.4 that $\Lambda^{(1)} = \Lambda^{(3)}$ on $\Sigma_1 \times (\delta_3, T_0 - \delta_3)$.

Denote by S_1 the union of all geodesics in $\overline{\Omega}^{(1)}$ with respect to the metric $\|g_1^{jk}\|^{-1}$, starting at $\hat{\Sigma}_1$, orthogonal to $\hat{\Sigma}_1$ and having lengths $\varepsilon > 0$. Since $\varepsilon > 0$ is small (we assume that $0 < \varepsilon \ll \delta_3 \ll \delta_2$) there is no focal points in $\overline{S}_1$ and the interior of S_1 does not intersect $\partial\Omega^{(1)}$. Since $\varepsilon \ll \delta_3$ we can apply Lemma 3.4 to $\overline{S}_1$ and $L^{(1)}, L^{(3)}$. We get that there is a diffeomorphism ψ_{41} of $\overline{S}_1$ onto $\overline{S}_1' \overset{def}{=} \psi_{41}(\overline{S}_1) \subset \overline{\Omega}^{(3)}$ and a gauge transformation $c_{41} \in G_0(\overline{S}_1)$, such that $\varphi_{41} = I$ on $\hat{\Sigma}_1$, $c_{41} = 1$ on $\hat{\Sigma}_1$ and $L^{(1)} = L^{(4)}$ on S_1, where $L^{(4)} \overset{def}{=} c_{41} \circ \psi_{41}^{-1} \circ L^{(3)}$.

Define $\psi_4' = \psi_{41}^{-1}$ on $\overline{S}_1'$, $\psi_4' = I$ on $\overline{R}_2$, $c_4' = c_{41}$ on $\overline{S}_1$, $c_4' = 1$ on $\overline{R}_2$. Since $L^{(1)} = L^{(3)}$ in $\overline{R}_2$ we have that ψ_4' is a C^∞ diffeomorphism on $\overline{S}_1' \cup \overline{R}_2$ and $c_4 \in G_0(\overline{S}_1 \cup \overline{R}_2)$ (c.f. Remark 2.3). Using Lemma 2.3 we can extend the diffeomorphism ψ_4' from $\overline{S}_1' \cup \overline{R}_2 \subset \overline{\Omega}^{(3)}$ to $\overline{\Omega}^{(3)}$. We can also extend c_4' from $\overline{S}_1 \cup \overline{R}_2$ to $\Omega^{(4)} \overset{def}{=} \psi_4'(\overline{\Omega}^{(3)})$.

Define $\psi_4 = \psi_4' \circ \psi_3$, $c_4 = c_4'(x)c_3(\psi_4^{-1}(x))$. Then $L^{(1)} = L^{(4)}$ in $\overline{S}_1 \cup \overline{R}_2$ and $L^{(4)} = c_4 \circ \psi_4 \circ L^{(2)}$ is an operator in $\Omega^{(4)}$.

Let σ_2 be the same as σ_1 with δ_3 replaced by $\delta_3 + \varepsilon$, let $\Sigma_2 = \psi_1^{-1}(\sigma_2)$ and let $\hat{\Sigma}_2$ be the smoothing of Σ_2 (see Fig.1). Denote by R_3 the domain bounded by $\hat{\Sigma}_2$ and $O(\delta_1 + \delta_2)$. Note that $R_2 \subset R_3 \subset \overline{S}_1 \cup \overline{R}_2$. Therefore $L^{(1)} = L^{(4)}$ in $\overline{R}_3$. It follows from Lemma 2.4 that $\Lambda^{(1)} = \Lambda^{(4)}$ on $\hat{\Sigma}_2 \times (\delta_2 + \varepsilon, T_0 - \delta_2 - \varepsilon)$.

Now repeat the same construction with $\hat{\Sigma}_1$ replaced by $\hat{\Sigma}_2$.

We will get a domain $S_2 \subset \overline{\Omega}^{(1)}$ consisting of all geodesics with respect to the metric $\|g_1^{jk}\|^{-1}$ starting on $\hat{\Sigma}_2$, orthogonal to $\hat{\Sigma}_2$ and having the length ε, where $\varepsilon \ll \delta_3$ is the same as above. Applying Lemma 3.4 and Remark 2.3 we get a diffeomorphism ψ_5' of $\overline{S}_2' \cup \overline{R}_3$ onto $\overline{S}_2 \cup \overline{R}_3$ and a gauge transformation $c_5' \in G_0(\overline{S}_2 \cup \overline{R}_3)$ such that $L^{(1)} = L^{(5)}$ on $\overline{S}_2 \cup \overline{R}_3$ where $L^{(5)} = c_5' \circ \psi_5' \circ L^{(4)}$, $S_2' \subset \overline{\Omega}^{(4)}$ is the same as S_2 with respect to the metric $\|g_4^{jk}\|^{-1}$. Using Lemma 2.3 we extend ψ_5' from $\overline{S}_2' \cup \overline{R}_3 \subset \overline{\Omega}^{(4)}$ to $\Omega^{(4)}$ as a diffeomorphism of $\overline{\Omega}^{(4)}$ onto $\overline{\Omega}^{(5)} \overset{def}{=} \psi_5'(\overline{\Omega}^{(4)})$. Define $\psi_5 = \psi_5' \circ \psi_4$, $c_5(x) = c_5'(x)c_4(\psi_5^{-1}(x))$. Then $\psi_5(x)$ is a diffeomorphism of $\overline{\Omega}^{(2)}$ onto $\overline{\Omega}^{(5)}$, $c_5(x) \in G_0(\overline{\Omega}^{(5)}$, $L^{(5)} = c_5 \circ \psi_5 \circ L^{(2)}$.

Analogously, let σ_3 be the same as σ_1 with δ_2 replaced by $\delta + 2\varepsilon$, $\Sigma_3 = \psi_1^{-1}(\sigma_3)$ and let $\hat{\Sigma}_3$ be a smoothing of Σ_3. Denote by R_4 the domain bounded by $\hat{\Sigma}_3$ and $O(\delta_1+\delta_2)$. Then $\overline{R}_3 \subset \overline{R}_4 \subset (\overline{S}_2 \cup \overline{R}_3)$ and $L^{(1)} = c_6 \circ \psi_6 \circ L^{(2)}$ on $\overline{R}_4$, where ψ_6 is a diffeomorphism of $\overline{\Omega}^{(2)}$ onto $\overline{\Omega}^{(6)} \overset{def}{=} \psi_6(\overline{\Omega}^{(2)})$ and $c_6 \in G_0(\overline{\Omega}^{(6)})$. After $\frac{T-\delta_3}{\varepsilon} = N$ steps we get a domain $\overline{R}_N \supset \overline{R}$, a diffeomorphism ψ_{N+2} of $\overline{\Omega}^{(2)}$ onto $\overline{\Omega}^{(N+2)}$ and a gauge transformation $c_{N+2} \in G_0(\overline{\Omega}^{(N+2)})$ such that $L^{(1)} = c_{N+2} \circ \psi_{N+2} \circ L^{(2)}$ on $\overline{R}_N$, $\psi_{N+2} = I$ and $c_{N+2} = 1$ on Γ_0.

We proved the following lemma:

Lemma 4.1. *Let $O' \subset \Gamma_0$ be a small neighborhood of $O(\delta_1 + \delta_2)$. Suppose $\Lambda^{(1)} = \Lambda^{(2)}$ on $O' \times (0, T_0)$. Let $R \subset \Omega^{(1)}$ be the same as above and $T < \frac{T_0}{2}$. Then there exists a diffeomorphism φ of R onto $\overline{R'} \overset{def}{=} \varphi(\overline{R}) \subset \overline{\Omega}^{(2)}$ and a gauge tranformation $c \in G_0(\overline{R})$ such that $L^{(1)} = c \circ \varphi^{-1} \circ L^{(2)}$ on $\overline{R}$, $\varphi = I$ and $c = 1$ on $O(\delta_1)$.*

Remark 4.1 In order to prove Lemma 4.1 we used $L^{(p)}, p = 1, 2$ only in a small neighborhoods of $\overline{R}$ and $\overline{R'}$. The properties of $L^{(p)}, p = 1, 2$, outside of these neighborhoods play no role. For example, $L^{(p)}, p = 1, 2$, are not required to be self-adjoint outside of neighborhoods of R and R'.

Remark 4.2 The following generalization of Lemma 4.1 holds:

Let $x^{(0)} \in \Gamma_0$ and let $x^{(1)}$ be an arbitrary point in $\Omega^{(1)}$. Let γ be an arbitrary curve in $\Omega^{(1)}$ connecting points $x^{(0)}$ and $x^{(1)}$ and having the length less than $\frac{T_0}{2}$. There exists a neighborhood $V_0 \subset \Omega^{(1)}$ of γ, a diffeomofphism ψ_0 of $\overline{V}_0$ onto $\overline{V}'_0 \overset{def}{=} \psi_0(\overline{V}_0) \subset \Omega^{(2)}$ and there exists a gauge $c_0(x) \in G_0(\overline{V}_0)$ such that $L^{(1)} = c_0 \circ \psi_0^{-1} \circ L^{(2)}$ in $\overline{V}_0$, $\psi_0 = I$ and $c_0 = 1$ on $\overline{V}_0 \cap \Gamma_0$ assuming that $\Lambda^{(1)} = \Lambda^{(2)}$ on $\tilde{O} \times (0, T_0)$ where $\tilde{O} \subset \Gamma_0$ is a small neighborhood of $\overline{V}_0 \cap \Gamma_0$.

To prove this result we approximate γ by a piece-wise smooth curve consisting of geodesic segments and prove Lemma 4.1 successively for each geodesic segment (see similsr arguments below). □

Now we actually start the global construction (c.f. [KKL] and [KKL1]). Since $\overline{\Omega}^{(1)}$ is compact there exists $\delta_0 > 0$ such that for any point $x^{(0)} \in \overline{\Omega}^{(1)}$ the geodesics starting at $x^{(0)}$ form a local system of coordinates in $B(x^{(0)}, \delta_0) \setminus B(x^{(0)}, \delta)$ for any $0 < \delta < \delta_0$. Here $B(x^{(0)}, r) = \{x : d(x, x^{(0)}) < r\}$ is a ball of radius r. We assume that the metric $\|g_1^{jk}\|^{-1}$ is extended to a δ_0-neighborhood of $\overline{\Omega}^{(1)}$. Therefore $B(x^{(0)}, \delta_0)$ make sense when $B(x^{(0)}, \delta_0) \cap \partial\Omega^{(1)} \neq \emptyset$. Let $\varepsilon_1 > 0$ be such that the semigeodesic coordinates with respect to $\partial\Omega^{(1)}$ hold in $2\varepsilon_1$-neighborhood of $\partial\Omega^{(1)}$ and the interior of this neighborhood does not intersect $\partial\Omega^{(1)}$. For each $x^{(0)} \in \partial\Omega^{(1)}$ consider a ball $B(x^{(0)}, \varepsilon_1)$. For each

$x^{(0)} \in \Omega^{(1)}$ consuder a ball $B(x^{(0)}, r)$ such that $r < \frac{\delta_0}{2}$ and $\overline{B}(x^{(0)}, r) \cap \partial\Omega^{(1)} = \emptyset$. Such balls form an open cover of $\overline{\Omega}^{(1)}$ and since $\overline{\Omega}^{(1)}$ is compact there exists a finite subcover $\{B(x^{(k)}, r_k)\}, k = 1, ..., N$. Denote by Ω_{ε_1} the union of all balls $B(x^{(k)}, r_k)$ such that $\overline{B}(x^{(k)}, r_k) \cap \partial\Omega^{(1)} = \emptyset$.

Let Γ and T be the same as in Lemma 2.2. Repeating the proof of Lemmas 2.2 and 2.3 and using th same notations we get a domain $B_1 \subset \Omega^{(1)}$, $\partial B_1 \cap \Gamma \neq \emptyset$ and connected, a diffeomorphism φ_3 of $\overline{\Omega}^{(2)}$ onto $\overline{\Omega}^{(3)} \overset{def}{=} \varphi_3(\overline{\Omega}^{(2)})$ and a gauge transformation $c_3 \in G_0(\overline{\Omega}^{(3)})$ such that $L^{(3)} = L^{(1)}$ in $B_1 \subset \Omega^{(1)} \cap \Omega^{(3)}$ where $L^{(3)} = c_3 \circ \varphi_3 \circ L^{(2)}$ is an operator in $\Omega^{(3)}$, $\varphi_3 = I$ on Γ_0. Since $\varphi_3 = I$ on Γ_0 we have that $\Lambda^{(3)} = \Lambda^{(1)}$ on $\Gamma_0 \times (0, T_0)$ where $\Lambda^{(3)}$ is the D-to-N for $L^{(3)}$. Let $\Gamma_1 = (\Gamma_0 \setminus (\Gamma_0 \cap \partial B_1)) \cup (\partial B_1 \setminus \Gamma_0)$. Note that $\Gamma_1 \subset \partial(\Omega^{(1)} \setminus \overline{B}_1) \cap \partial(\Omega^{(3)} \setminus \overline{B}_1)$.



Figure 2. The domain U_0. The boundary S_0 is drawn bold.

Take arbitrary point $x^{(0)} \in B_1$ such that $d(x^{(0)}, \Gamma_1) = 2\varepsilon_2$ where ε_2 is much smaller than δ_0. We have that $B(x^{(0)}, \varepsilon_2) \subset B_1$. Pick some geodesics γ_0 starting at $x^{(0)}$ and denote by U_0 the part of $B(x^{(0)}, \delta) \setminus \overline{B}(x^{(0)}, \varepsilon_2)$ consisting of all geodesics γ starting at $x^{(0)}$ and having an angle less than $\pi - \varepsilon_3$ with γ_0 at $x^{(0)}$. We choose $\delta \leq \delta_0, \gamma_0$ and ε_3 such that $\overline{U}_0 \cap \partial\Omega^{(1)} = \emptyset$. Denote by S_0 a smooth surface in B_1 that contains $\overline{U}_0 \cap \partial B(x^{(0)}, \varepsilon_2)$ and divides B_1 in two domains, B_1^+ and B_1^-, where B_1^+ is bounded by Γ_1 and S_0, and $B_1^- = B_1 \setminus \overline{B}_1^+$ (see Fig.2). We assume that $\Omega^{(1)} \setminus \overline{B}_1^-$ has a smooth boundary that includes S_0. Denote $\Gamma_2 = S_0 \cup (\Gamma_1 \setminus \partial B_1^+)$. Since $L^{(3)} = L^{(1)}$ in $\overline{B}_1^-$ and $\Lambda^{(3)} = \Lambda^{(1)}$ on $\Gamma_0 \times (0, T)$ we get from the Lemma 2.4 that $\Lambda^{(3)} = \Lambda^{(1)}$ on $\Gamma_2 \times (\delta', T - \delta')$, where $\delta' = \max_{x \in \bar{B}_1^-} d_1(x, \Gamma_0)$ and we consider the D-to-N operators corresponding to $L^{(1)}, L^{(3)}$ in domains $(\Omega^{(1)} \setminus \overline{B}_1^-) \times (\delta', T_0 - \delta')$, $(\Omega^{(3)} \setminus \overline{B}_1^-) \times (\delta', T_0 - \delta')$.

Now apply Lemma 4.1 to $U_0 \subset (\Omega^{(1)} \setminus B_1^-)$ instead of R. Let U_0' be the union of all geodesics in $\Omega^{(3)}$ with respect to the metric $\|g_3^{jk}\|^{-1}$ starting at $\overline{U}_0 \cap \partial B(x^{(0)}, \varepsilon_2)$, orthogonal to $\partial B(x^{(0)}, \varepsilon_2)$ and having the length $\delta - \varepsilon_2$. Note that $B(x^{(0)}, \varepsilon_2) \subset B_1 \subset \Omega^{(1)} \cap \Omega^{(3)}$. Since $\Lambda^{(1)} = \Lambda^{(3)}$ on $\Gamma_2 \times (\delta', T_0 - \delta')$, the Lemma 4.1 implies that there exists a diffeomorphism φ of $\overline{U}_0' \subset \overline{\Omega}^{(2)}$ onto $\overline{U}_0$ and there exists $c(x) \in C^\infty(\overline{U}_0)$, $c(x) \neq 0$ in $\overline{U}_0$ such that $c \circ \varphi \circ L^{(3)} = L^{(1)}$ in $\overline{U}_0$, $\varphi = I$ and $c = 1$ on $S_0 \cap \overline{U}_0$. Define $\varphi = I$ on $\overline{B}_1 \setminus U_0$ and Γ_0. Also define $c = 1$ on $\overline{B}_1 \setminus U_0$ and Γ_0. Since $L^{(3)} = L^{(1)}$ in $\overline{U}_0 \cap \overline{B}_1$ we have that $\varphi = I$ and $c = 1$ in $\overline{U}_0 \cap \overline{B}_1$. Therefore φ and c are C^∞ on $\overline{U}_0 \cup \overline{B}_1 \cup \Gamma_0$ (c.f. Remark 2.3). Applying Lemma 2.3 we can extend φ from $\overline{U}_0 \cup \overline{B}_1 \cup \Gamma_0$ to $\overline{\Omega}^{(3)}$ as a diffeomorphism of $\overline{\Omega}^{(3)}$ onto $\overline{\Omega}^{(4)} \overset{def}{=} \varphi(\overline{\Omega}^{(3)})$ and extend c from $\overline{U}_0 \cup \overline{B}_1 \cup \Gamma_0$ to $\overline{\Omega}^{(4)}$ as an element of $G_0(\overline{\Omega}^{(4)})$.

Let $\varphi_4 = \varphi \circ \varphi_3$. Then φ_4 is a diffeomorphism of $\overline{\Omega}^{(2)}$ onto $\overline{\Omega}^{(4)}$. Let $c_4(x) = c(x) c_3(\varphi_4^{-1}(x)) \in G_0(\Omega^{(4)}$. Then we have $L^{(4)} = c_4 \circ \varphi_4 \circ L^{(2)}$ is an operator in $\Omega^{(4)}$ equal to $L^{(1)}$ in $\overline{U}_0 \cup \overline{B}_1 \subset \overline{\Omega}^{(1)} \cap \overline{\Omega}^{(4)}$.

Therefore applying Lemma 4.1 to U_0 we gained that $\overline{B}_1$ is replaced by a larger domain $\overline{U}_0 \cup \overline{B}_1$.

Taking a point $x^{(1)} \in U_0 \cup B_1$ instead of $x^{(0)}$ we can construct a domain U_1 similar to U_0. For the brevity we shall call by U-type domains the domains similar to U_0. We shall show that adding a finite number of U-type domains we can cover $\overline{\Omega}^{(1)}$.

Take any ball $B(x^{(p)}, r_p) \subset \Omega_{\varepsilon_1}$. Since $\overline{\Omega}^{(1)}$ is connected, the point $x^{(p)}$ can be connected with $x^{(0)} \in B_1$ by a broken geodesics (c.f. [KKL]), more exactly, there exist points $y_1 = x^{(0)}, y_2, ..., y_{N_1} = x^{(p)}$ such that y_k and $y_{k+1}, 1 \leq k \leq N_1 - 1$, can be connected by a geodesics of length $\leq \frac{\delta_0}{2}$. Using a sequence of U-type domains we can cover this broken geodesics including $x^{(p)}$. Adding more U-type domains if needed we can cover $B(x^{(p)}, r_p)$ too. We can do this with any ball $B(x^{(p)}, r_p) \subset \Omega_{\varepsilon_1}$. Therefore inserting M of U-type domains

$U_1, ..., U_M$ in $\Omega^{(1)}$ we get a sequences of domains $\Omega^{(k)}, 4 \le k \le M+4$, diffeomorphisms φ_k of $\overline{\Omega}^{(2)}$ onto $\overline{\Omega}^{(k)}$ and gauge transformations $c_k \in G_0(\overline{\Omega}^{(k)})$ such that $L^{1)} = L^{(k)}$ in $\overline{B}_1 \cup (\cup_{j=0}^{k-4} \overline{U}_j),\ 4 \le k \le M+4$, where $L^{(k)} = c_k \circ \varphi_k \circ L^{(2)}$ are operators in $\Omega^{(k)},\ B_1 \cup (\cup_{j=0}^{k-4} U_j) \subset \Omega^{(1)} \cap \Omega^{(k)},\ \Lambda^{(1)} = \Lambda^{(k)}$ on $\Gamma_0 \times (0, T_0)\ \ 4 \le k \le M+4$.

Let M be such that $\Omega_{\varepsilon_1} \subset B_1 \cup (\cup_{j=0}^{M} U_j) \subset \Omega^{(1)} \cap \Omega^{(M+4)}$. We have $L^{(1)} = L^{(M+4)}$ in $\overline{\Omega}_{\varepsilon_1}$, and $\Lambda^{(1)} = \Lambda^{(M+4)}$ on $\Gamma_0 \times (0, T_0)$.

Note that we choose $U_j, 0 \le j \le M$, taking into account the geometry of $\Omega^{(1)}$ and the metric $\|g_1^{jk}\|^{-1}$, regardless of the geometry of $\Omega^{(p)}$ and the metric $\|g_p^{jk}\|^{-1},\ 2 \le p$.

Now consider the cover of $\overline{\Omega}^{(1)} \setminus \Omega_{\varepsilon_1}$ by U-type domains. Let $\alpha = \partial\Omega^{(1)} \cap B(x^{(k)}, \varepsilon_1)$, where $\{B(x^{(k)}, \varepsilon_1)\}$ is a finite cover of $\partial\Omega^{(1)}$. Denote by U_{M+1} the union of all geodesics with respect to the metric $\|g_1^{jk}\|^{-1}$ starting on $\overline{\alpha}$, orthogonal to $\overline{\alpha}$ and having the same lengths $2\varepsilon_1$. Let W_{M+1} be the set of endpoints of these geodesics. It follows from the definition of Ω_{ε_1} that W_{M+1} is located inside Ω_{ε_1}.

Denote by U'_{M+1} the union of all geodesics with respect to metric $\|g_{M+4}^{jk}\|^{-1}$ corresponding to $L^{(M+4)}$ that start on W_{M+1}, orthogonal to W_{M+1} and having the lengths $2\varepsilon_1$. Note that $W_{M+1} \subset \Omega_{\varepsilon_1} \subset \Omega^{(M+4)}$.

Let $\tilde{W}_{M+1}$ bea surface in Ω_{ε_1} containing W_{M+1} and similar to S_0 in the case of domain U_0. Since $\Lambda^{(1)} = \Lambda^{(M+4)}$ on $\Gamma_0 \times (0, T_0)$ and $L^{(1)} = L^{(M+4)}$ on $\overline{\Omega}_{\varepsilon_1}$, Lemma 2.4 implies that $\Lambda^{(1)} = \Lambda^{(M+4)}$ on $\tilde{W}_{M+4} \times (T', T_0 - T')$, where $T' = \max_{x \in \overline{\Omega}_{\varepsilon_1}} d_1(x, \Gamma_0)$. Now we can repeat for U_{M+1} the same arguments as for U_0. Applying Lemma 4.1 we get that there exists a diffeomorphism φ'_{M+5} of $\overline{U}'_{M+1}$ onto $\overline{U}_{M+1},\ \varphi'_{M+5} = I$ on W_{M+1} and $c'_{M+5}(x) \in G_0(\overline{U}_{M+1}),\ c' = 1$ on W_{M+1} such that

$$c'_{M+5} \circ \varphi'_{M+5} \circ L^{(M+4)} = L^{(1)} \text{ in } U_{M+1}. \tag{4.1}$$

Since $L^{(M+4)} = L^{(1)}$ in Ω_{ε_1} we have that $\varphi'_{M+5} = I$ on $\overline{U}_{M+5} \cap \overline{\Omega}_{\varepsilon_1}$. Therefore φ'_{M+5} is a diffeomorphism on $\overline{U}_{M+5} \cup \overline{\Omega}_{\varepsilon_1}$ after extending φ'_{M+5} as I on $\overline{\Omega}_{\varepsilon_1}$ (c.f. Remark 2.3). Analoguously taking $c'_{M+5} = 1$ on $\overline{\Omega}_{\varepsilon_1}$ we get a C^∞-function on $\overline{\Omega}_{\varepsilon_1} \cup \overline{U}_{M+1}$. Note that Lemma 4.1 implies that $U'_{M+1} \subset \Omega^{(M+4)}$.

We shall consider first the case when $\alpha \cap \Gamma_0 \neq \emptyset$. We shall prove that $\varphi'_{M+5} = I,\ c'_{M+5} = 1$ on $\alpha_1 \overset{def}{=} \alpha \cap \Gamma_0$.

Let g be arbitrary smooth function with the support in $\overline{\alpha}_1 \times (T', T_0 - T')$.

Let u_1 be the solution of $L_1 u_1 = 0$ on $\Omega^{(1)} \times (0, T_0),\ u_1 = u_{1t} = 0$ for $t = 0,\ u_1|_{\Gamma_0 \times (0, T_0)} = g$. Also let $u_{M+4}(x)$ be the solution of $L^{(M+4)} u_{M+4} = 0$

in $\Omega^{(M+4)}\times(0,T_0)$, $u_{M+4}=\frac{\partial u_{M+4}}{\partial t}=0$ for $t=0$ and $u_{M+4}|_{\Gamma_0\times(0,T_0)}=g$. Since $\Lambda^{(1)}=\Lambda^{(M+4)}$ on $\Gamma_0\times(0,T)$ and since $L^{(M+4)}=L^{(1)}$ in $\overline{\Omega}_{\varepsilon_1}$ we get, by the unique continuation theorem (c.f [T]), that $u_1=u_{M+4}$ in $\overline{\Omega}_{\varepsilon_1}\times(T',T_0-T')$.

It follows from (4.1) that $u_1(x,t)$ and $u_{M+5}(x,t)=c'_{M+5}(x)u_{M+4}((\varphi'_{M+5})^{-1}(x),t)$ satisfy the same equation $L^{(1)}u=0$ in $U_{M+1}\times(T',T_0-T'))$. Since $u_1(x)=u_{M+4}(x)$, $\varphi'_{M+5}=I$ and $c'_{M+5}=1$ on $\Omega_{\varepsilon_1}\times(T',T_0-T')$ we have that u_1 and u_{M+5} have the same Cauchy data on $\widetilde{W}_{M+1}\times(T',T_0-T')$.

We have, using again the unique continuation theorem that

$$c'_{M+5}(x)u_{M+4}((\varphi'_{M+5})^{-1}(x),t)=u_1(x,t) \text{ in } \overline{U}_{M+1}\times(T_1+2\varepsilon_1,T_0-T_1-2\varepsilon_1). \tag{4.2}$$

Taking the restriction of (4.2) to $\overline{\alpha}_1\times(T'+2\varepsilon_1,T_0-T'-2\varepsilon_1)$ we obtain

$$c'_{M+5}(x)g((\varphi'_{M+5})^{-1}(x),t)=g(x,t) \text{ in } \overline{\alpha}_1\times(T_1+2\varepsilon_1,T_0-T_1-2\varepsilon_1). \tag{4.3}$$

Since g is arbitrary we get that $\varphi'_{M+5}=I$ on $\overline{\alpha}_1$, $c'_{M+5}(x)=1$ on $\overline{\alpha}_1$. Therefore, we take $\varphi'_{M+5}=I$, $c'_{M+1}(x)=1$ on Γ_0 and get a C^∞-functions on $\Gamma_0\cup\overline{\Omega}_{\varepsilon_1}\cup\overline{U}_{M+1}$. There is no any obstruction to the extension $\varphi'_{M+5}=I$, $c'_{M+5}=1$ on Γ_0 in the case when $\alpha\cap\Gamma_0=\emptyset$. Thus in both cases applying Lemma 2.3 we get a diffeomorphism φ'_{M+5} of $\Omega^{(M+4)}$ onto $\overline{\Omega}^{(M+5)}\overset{def}{=}\varphi'_{M+5}(\overline{\Omega}^{(M+4)})$ and $c'_{M+5}\in G_0(\Omega^{(M+5)})$ such that $\varphi'_{M+5}=I$ on Γ_0.

As in the case of the U-type domain U_0, denote $\varphi_{M+5}=\varphi'_{M+5}\circ\varphi_{M+4}$, $c_{M+5}=c'_{M+5}c_{M+4}(\varphi^{-1}_{M+4}(x))$. Then we get that

$$L^{(M+5)}=L^{(1)} \quad \text{in} \quad \overline{\Omega}_{\varepsilon_1}\cup\overline{U}_{M+1}\subset\overline{\Omega}^{(1)}\cap\overline{\Omega}^{(M+5)},$$

where $L^{(M+5)}=c_{M+5}\circ\varphi_{M+5}\circ L^{(2)}$ is an operator on $\overline{\Omega}^{(M+5)}$, φ_{M+5} is a diffeomorphism of $\overline{\Omega}^{(2)}$ onto $\overline{\Omega}^{(M+5)}$, $\varphi_{M+5}=I$ on Γ_0, $c_{M+5}\in G_0(\overline{\Omega}^{(M+5)})$. In particular, we have $\alpha\subset\partial\Omega^{(1)}\cap\partial\Omega^{(M+5)}$ and φ^{-1}_{M+5} maps $\alpha\subset\partial\Omega^{(1)}$ onto $\alpha'\in\partial\Omega^{(2)}$. Repeating the same construction with each of $B(x^{(k)},\varepsilon_1)$ such that $B(x^{(k)},\varepsilon_1)\cap\partial\Omega^{(1)}\neq\emptyset$ we get $\overline{\Omega}_{\varepsilon_1}\cup(\cup_{j=1}^{M_1}\overline{U}_{M+j})=\overline{\Omega}^{(1)}$, $\overline{\Omega}_{\varepsilon_1}\cup(\cup_{j=1}^{M_1}\overline{U}_{M+j})\subset\overline{\Omega}^{(M+M_1+4)}$ and $L^{(1)}=c_{M+M_1+4}\circ\varphi_{M+M_1+4}\circ L^{(2)}$ in $\overline{\Omega}_{\varepsilon_1}\cup(\cup_{j=1}^{M_1}\overline{U}_j)$, where $\overline{\Omega}^{M+M_1+4)}\overset{def}{=}\varphi_{M+M_1+4}(\overline{\Omega}^{(2)})$, $\varphi_{M+M_1+4}(x)$ is a diffeomorphism, $\varphi_{M+M_1+4}=I$ on Γ_0, $c_{M+M_1+4}\in G_0(\Omega^{M+M_1+4)})$.

We claim that $\Omega^{(M+M_1+4)}=\Omega^{(1)}$. If $\overline{\Omega}^{(M+M_1+4)}\neq\overline{\Omega}^{(1)}$ then there exists an interior point of $\Omega^{(M+M_1+4)}$ that is a boundary point of $\Omega^{(1)}$. It follows from the arguments similar to the proof of Lemma 3.2 that this is impossible. □

References

[B1] Belishev, M., 1997, Boundary control in reconstruction of manifolds and metrics (the BC method), Inverse Problems 13, R1-R45

[B2] Belishev, M., 2002, How to see waves under the Earthsurface (the BC-method for geophysicists), Ill -Posed and Inverse Problems, 55-72 (S.Kabanikhin and V.Romanov (Eds), VSP)

[E] Eskin, G. 2006 , A new approach to hyperbolic inverse problems, Inverse Problems 22, 815-833

[E1] Eskin, G., 2004, Inverse boundary value problems in domains with several obstacles, Inverse Problems 20, 1497-1516

[E2] Eskin, G., 2005, Inverse problems for Schrödinger equations with Yang-Mills potentials in domains with obstacles and the Aharonov-Bohm effect, Journal of Physics Conference Series, 12, 23-32

[Hi] Hirsch, M., 1976, Differential topology (New York : Springer)

[I] Isakov, V., 1998, Inverse problems for partial differential equations, Appl. Math. Studies, vol. 127, Springer, 284 pp.

[K] Kurylev, Y., 1993, Multi-dimensional inverse boundary problems by BC-mathod : groups of transformations and uniqueness results, Math. Comput. Modelling 18, 33-45

[KK] Katchalov, A., Kurylev, Y., 1998, Multidimensional inverse problems with incomplete boundary spectral data, Comm. Part. Diff. Eq. 23, 55-95

[KKL] Katchalov, A., Kurylev, Y., Lassas, M., 2001, Inverse boundary spectral problems (Boca Baton : Chapman&Hall)

[KKL1] Katchalov, A., Kurylev, Y., Lassas, M., 2004, Energy measurements and equivalence of boundary data for inverse problems on noncompact manifolds, IMA Volumes, v.137, 183-214

[KL1] Kurylev, Y. and Lassas, M., 2000, Hyperbolic inverse problems with data on a part of the boundary AMS/1P Stud. Adv. Math, 16, 259-272

[T] Tataru, D., 1995, Unique continuation for solutions to PDE, Comm. in PDE 20, 855-84